Neuroscience Letters, 187 (1995) 215–218
Elsevier Science Ireland Ltd.

AUTHOR INDEX

Adrio, F., see Alvarez-Otero, R., 87
Agren, G., Lundeberg, T., Uvnäs-Moberg, K., Sato, A., The oxytocin antagonist 1-deamino-2-D-Tyr-(Oet)-4-Thr-8-Orn-oxytocin reverses the increase in the withdrawal response latency to thermal, but not mechanical nociceptive stimuli following oxytocin administration or massage-like stroking in rats, 49
Alonso Solís, R., see Benítez, L., 37
Alvarez-Otero, R., Pérez, S.E., Rodríguez, M.A., Adrio, F., Anadón, R., GABAergic neuronal circuits in the cerebellum of the dogfish *Scyliorhinus canicula* (Elasmobranchs): an immunocytochemical study, 87
Anadón, R., see Alvarez-Otero, R., 87
Andersen, S., see Ebert, B., 165
Anderson-Beck, R., Wilson, L., Brazier, S., Hughes, I.E., Peers, C., Doxapram stimulates dopamine release from the intact rat carotid body in vitro, 25
Armstrong, R.A., Myers, D., Smith C.U.M., What determines the size frequency distribution of β-amyloid (Aβ) deposits in Alzheimer's disease patients? 13
Artigas, F., see Ferré, S., 61
Asano, T., see Otsuguro, K., 145

Balderson, D.J., see Pickering-Brown, S.M., 127
Bartfai, T., see Bristulf, J., 53
Bartfai, T., see Valkna, A., 75
Beckmann, H., see Künig, G., 107
Benítez, L., Fanjul, L.F., Ruiz de Galarreta, C.M., Quintana Aguiar, J., González Reyes, J., Hernández, I., Santana Delgado, P., Cabrera Oliva, J., Alonso Solís, R., Estévez Rosas, F., Partial characterization of a thyrotropin releasing hormone-sensitive glycosylphosphatidylinositol in pituitary lactotrophes, 37
Benoist, J.-M., see Guilbaud, G., 197
Birtsch, C., see Schröder, H., 173
Block, G.D. see Khalsa, S.B.S., 41
Bonkale, W.L., Winblad, B., Ravid, R., Cowburn, R.F., Reduced nitric oxide responsive soluble guanylyl cyclase activity in the superior temporal cortex of patients with Alzheimer's disease, 5
Bonnard, M., see Danion, F., 185
Brandenberger, G., see Weibel, L., 83
Brazier, S., see Anderson-Beck, R., 25
Bristulf, J., Bartfai, T., Interleukin-1β and tumour necrosis factor-α stimulate the mRNA expression of interleukin-1 receptors in mouse anterior pituitary AtT-20 cells, 53
Bromm, B., Scharein, E., Darsow, U., Ring, J., Effects of menthol and cold on histamine-induced itch and skin reactions in man, 157

Cabrera Oliva, J., see Benítez, L., 37
Charness, M.E., see Perides, G., 21
Cherubini, E., see Sciancalepore, M., 91
Chu, L., Kaneko, C.R.S., Do the saccadic system and the vestibulo-ocular reflex of monkeys share a common neural integrator? 193
Chung, S.-H., Kuyucak, S., Changes in the kinetics and conductance of *N*-methyl-D-aspartate (NMDA)-receptor activated single channels with temperature, 181
Cooper, A.J., Moser, B., Mitchell, I.J., A subset of striatopallidal neurons are Fos-immunopositive following acute monoamine depletion in the rat, 189
Cooper, P.N., see Pickering-Brown, S.M., 127
Cowburn, R.F., see Bonkale, W.L., 5
Craft, C.M., see Gauer, F., 69
Cruwys, S.C., see Garrett, N.E., 201

Danion, F., Bonnard, M., Pailhous, J., Joint-dependent mechanisms to adapt to an imbalance between flexion and extension forces in human gait, 185
Darsow, U., see Bromm, B., 157
de Silva, R., see Pickering-Brown, S.M., 127
de Vos, R.A.I., see Schröder, H., 173
Deckert, J., see Künig, G., 107
Deusch, G., see Glocker, F.X., 161

Ebert, B., Andersen, S., Krogsgaard-Larsen, P., Ketobemidone, methadone and pethidine are non-competitive *N*-methyl-D-aspartate (NMDA) antagonists in the rat cortex and spinal cord, 165
Edgecomb, P., see Perides, G., 21
Ehrhart, J., see Weibel, L., 83
Eisenberg, E., LaCross, S., Strassman, A.M., The clinically tested *N*-methyl-D-aspartate receptor antagonist memantine blocks and reverses thermal hyperalgesia in a rat model of painful mononeuropathy, 17
Estévez Rosas, F., see Benítez, L., 37

Fanjul, L.F., see Benítez, L., 37
Ferré, S., Artigas, F., Clozapine decreases serotonin extracellular levels in the nucleus accumbens by a dopamine receptor-independent mechanism, 61
Follenius, M., see Weibel, L., 83
Foster, R.G., see García-Fernández, J.M., 33
Fu, Q.-G., see Helmchen, C., 137
Fujii, S., Ito, K.-I., Osada, H., Hamaguchi, T., Kuroda, Y., Kato, H., Extracellular phosphorylation of membrane protein modifies theta burst-induced long-term potentiation in CA1 neurons of guinea-pig hippocampal slices, 133
Fujii, S., Kato, H., Furuse, H., Ito, K.-I., Osada, H., Hamaguchi, T., Kuroda, Y., The mechanism of ATP-induced long-term potentiation involves extracellular phosphorylation of membrane proteins in guinea-pig hippocampal CA1 neurons, 130
Funada, M., see Suzuki, T., 45
Furukawa, H., see Nakagawa, Y., 209
Furuse, H., see Fujii, S., 130

Galteau, M.-M., see Jolivalt, C., 65
García-Fernández, J.M., Jimenez, A.J., Foster, R.G., The persistence of cone photoreceptors within the dorsal retina of aged retinally degenerate mice (*rd/rd*): implications for circadian organization, 33
Garrett, N.E., Kidd, B.L., Cruwys, S.C., Tomlinson, D.R., Streptozotocin-induced diabetes decreases substance P levels in experimental arthritis in the rat knee, 201
Gauer, F., Kedzierski, W., Craft, C.M., Identification of circadian gene expression in the rat pineal gland and retina by mRNA differential display, 69
Gautron, M., see Guilbaud, G., 197
Glocker, F.X., Guschlbauer, B., Lücking, C.H., Deusch, G., Effects of local injections of botulinum toxin on electrophysiological parameters in patients with hemifacial spasm: role of synaptic activity and size of motor units, 161

Goichot, B., see Weibel, L., 83
González Reyes, J., see Benítez, L., 37
Gross, D.R., see Sparks, D.L, 142
Guilbaud, G., Benoist, J.-M., Gautron, M., Contribution of the sciatic and saphenous nerve to the ventrobasal thalamic neuronal responses to pinch in rats with a chronic sciatic nerve constriction: a study using anesthetic blocks and nerve section, 197
Guschlbauer, B., see Glocker, F.X., 161

Hamaguchi, T., see Fujii, S., 130
Hamaguchi, T., see Fujii, S., 133
Hartmann, J., see Künig, G., 107
Heinsen, H., see Künig, G., 107
Helisalmi, S., see Soininen, H., 79
Helmchen, C., Fu, Q.-G., Sandkühler, J., Inhibition of spinal nociceptive neurons by microinjections of somatostatin into the nucleus raphe magnus and the midbrain periaqueductal gray of the anesthetized cat, 137
Hernández, I., see Benítez, L., 37
Hirata, K., Ohno-Shosaku, T., Sawada, S., Yamamoto, C., Baclofen inhibits GABAergic transmission after treatment with type-specific calcium channel blockers in cultured rat hippocampal neurons, 205
Hosaka, N., see Inouye, T., 29
Hughes, I.E., see Anderson-Beck, R., 25

Inoue, K., see Koizumi, S., 123
Inouye, T., Shinosaki, K., Toi, S., Matsumoto, Y., Hosaka, N., Potential flow of α activity in the human electroence phalogram, 29
Ironside, J.W., see Pickering-Brown, S.M., 127
Ito, K.-I., see Fujii, S., 130
Ito, K.-I., see Fujii, S., 133
Ito, S., see Otsuguro, K., 145

Jansen, E.N.H., see Schröder, H., 173
Jensen, F.E., see Perides, G., 21
Jimenez, A.J., see García-Fernández, J.M., 33
Jolivalt, C., Minn, A., Vincent-Viry, M., Galteau, M.-M., Siest, G., Dextromethorphan *O*-demethylase activity in rat brain microsomes, 65
Juréus, A., see Valkna, A., 75

Kaneko, C.R.S., see Chu, L., 193
Karelson, E., see Valkna, A., 75
Karhunen, T., Tilgmann, C., Ulmanen, I., Panula, P., Catechol-*O*-methyltransferase (COMT) in rat brain: immunoelectron microscopic study with an antiserum against rat recombinant COMT protein, 57
Kataoka, Y., see Koizumi, S., 123
Kato, H., see Fujii, S., 130
Kato, H., see Fujii, S., 133
Kedzierski, W., see Gauer, F., 69
Khalsa, S.B.S., Michel, S., Block, G.D., The anesthetic agents pentobarbital and chloralose block phase shifts of a neuronal in vitro circadian pacemaker, 41
Kidd, B.L., see Garrett, N.E., 201
Kitamoto, T., see Nakagawa, Y., 209
Kohzuma, M., see Koizumi, S., 123
Koizumi, S., Kataoka, Y., Inoue, K., Kohzuma, M., Niwa, M., Taniyama, K., Contribution of L-type Ca^{2+} channels to long-term enhancement of high K^+-evoked release of dopamine from rat striatal slices, 123
Koob, G.F., see Richter, R.M., 169
Kosunen, O., see Soininen, H., 79
Krause, F., see Künig, G., 107
Krogsgaard-Larsen, P., see Ebert, B., 165
Künig, G., Hartmann, J., Krause, F., Deckert, J., Heinsen, H., Ransmayr, G., Beckmann, H., Riederer, P., Regional differences in the interaction of the excitotoxins domoate and L-β-oxalyl-aminoalanine with [^{3}H]kainate binding sites in human hippocampus, 107
Kuroda, Y., see Fujii, S., 130
Kuroda, Y., see Fujii, S., 133
Kuyucak, S., see Chung, S.-H., 181

LaCross, S., see Eisenberg, E., 17
Lang, B, see Martin-Moutot, N., 115
Langel, Ü., see Valkna, A., 75
León-Olea, M., see Talavera, E., 177
Liu, H., see Sparks, D.L, 142
Lobron, C., see Schröder, H., 173
Lücking, C.H., see Glocker, F.X., 161
Lundeberg, T., see Agren, G., 49

Maeda, J., see Suzuki, T., 45
Maelicke, A., see Schröder, H., 173
Mann, D.M.A., see Pickering-Brown, S.M., 127
Mannermaa, A., see Soininen, H., 79
Martin-Moutot, N., Lang, B, Newsom-Davis, J, Seagar, M., Binding of Lambert-Eaton myasthenic syndrome IgG to synaptosomal proteins does not correlate with an inhibition of calcium uptake, 115
Martínez-Lorenzana, G., see Talavera, E., 177
Mashimo, T., see Shibuta, S., 103
Matsumoto, Y., see Inouye, T., 29
Meßlinger, K., see Schepelmann, K., 212
Michel, S. see Khalsa, S.B.S., 41
Minn, A., see Jolivalt, C., 65
Misawa, M., see Suzuki, T., 45
Mitchell, I.J., see Cooper, A.J., 189
Moser, B., see Cooper, A.J., 189
Mulvaney, J.M., Parsons, R.L., Arachidonic acid may mediate the galanin-induced hyperpolarization in parasympathetic neurons from *Necturus maculosus,* 95
Myers, D., see Armstrong, R.A., 13

Nakagawa, Y., Kitamoto, T., Furukawa, H., Ogomori, K., Tateishi, J., Allelic variation of apolipoprotein E in Japanese sporadic Creutzfeldt-Jakob disease patients, 209
Nakazato, Y., see Otsuguro, K., 145
Newsom-Davis, J, see Martin-Moutot, N., 115
Niwa, M., see Koizumi, S., 123
Nowacki, S., see Schröder, H., 173

Ogomori, K., see Nakagawa, Y., 209
Ohara, A., see Shibuta, S., 103
Ohno-Shosaku, T., see Hirata, K., 205
Ohta, T., see Otsuguro, K., 145
Osada, H., see Fujii, S., 130
Osada, H., see Fujii, S., 133
Otsuguro, K., Asano, T., Ohta, T., Ito, S., Nakazato, Y., ATP-evoked membrane current in guinea pig adrenal chromaffin cells, 145
Owen, F., see Pickering-Brown, S.M., 127

Pai, Y.-C., see Rogers, M.W., 99

Pailhous, J., see Danion, F., 185
Paljärvi, L., see Soininen, H., 79
Panula, P., see Karhunen, T., 57
Parsons, R.L., see Mulvaney, J.M., 95
Peers, C., see Anderson-Beck, R., 25
Pellicer, F., see Talavera, E., 177
Pérez, S.E., see Alvarez-Otero, R., 87
Perides, G., Jensen, F.E., Edgecomb, P., Rueger, D.C., Charness, M.E., Neuroprotective effect of human osteogenic protein-1 in a rat model of cerebral hypoxia/ischemia, 21
Pich, E.M., see Richter, R.M., 169
Pickering-Brown, S.M., Mann, D.M.A., Owen, F., Ironside, J.W., de Silva, R., Roberts, D.A., Balderson, D.J., Cooper, P.N., Allelic variations in apolipoprotein E and prion protein genotype related to plaque formation and age of onset in sporadic Creutzfeldt-Jakob disease, 127

Quintana Aguiar, J., see Benítez, L., 37

Ransmayr, G., see König, G., 107
Ravid, R., see Bonkale, W.L., 5
Reynolds, G.P., see Sardar, A.M., 9
Rice, A.S.C., Topical spinal administration of a nitric oxide synthase inhibitor prevents the hyper-reflexia associated with a rat model of persistent visceral pain, 111
Richter, R.M., Pich, E.M., Koob, G.F., Weiss, F., Sensitization of cocaine-stimulated increase in extracellular levels of corticotropin-releasing factor from the rat amygdala after repeated administration as determined by intracranial microdialysis, 169
Riederer, P., see König, G., 107
Riekkinen Sr, P., see Soininen, H., 79
Ring, J., see Bromm, B., 157
Roberts, D.A., see Pickering-Brown, S.M., 127
Rodríguez, M.A., see Alvarez-Otero, R., 87
Rogers, M.W., Pai, Y.-C., Organization of preparatory postural responses for the initiation of lateral body motion during goal directed leg movements, 99
Rueger, D.C., see Perides, G., 21
Ruiz de Galarreta, C.M., see Benítez, L., 37
Ryynänen, M., see Soininen, H., 79

Sánchez -Alvarez, M., see Talavera, E., 177
Sánchez-Islas, E., see Talavera, E., 177
Sandkühler, J., see Helmchen, C., 137
Santana Delgado, P., see Benítez, L., 37
Sardar, A.M., Reynolds, G.P., Frontal cortex indoleamine-2,3-dioxygenase activity is increased in HIV-1-associated dementia, 9
Sato, A., see Agren, G., 49
Sawada, S., see Hirata, K., 205
Schäfer, F., see Schmid, H.A., 149
Schaible, H.-G., see Schepelmann, K., 212
Scharein, E., see Bromm, B., 157
Scheff, S.W., see Sparks, D.L, 142
Schepelmann, K., Meßlinger, K., Schaible, H.-G., Schmidt, R.F., The opioid antagonist naloxone does not alter discharges of nociceptive afferents from the acutely inflamed knee joint of the cat, 212
Schmid, H.A., Schäfer, F., Simon E., Opposite effects of angiotensin II and nitric oxide on neurons in the duck subfornical organ, 149
Schmidt, R.F., see Schepelmann, K., 212
Schröder, H., de Vos, R.A.I., Jansen, E.N.H., Birtsch, C., Wevers, A., Lobron, C., Nowacki, S., Schröder, R., Maelicke, A., Gene expression of the nicotinic acetylcholine receptor $\alpha 4$ subunit in the frontal cortex in Parkinson's disease patients, 173
Schröder, R., see Schröder, H., 173
Sciancalepore, M., Cherubini, E., Protein kinase A-dependent increase in frequency of miniature GABAergic currents in rat CA3 hippocampal neurons, 91
Seagar, M., see Martin-Moutot, N., 115
Shamir, D., see Yanai, J., 153
Shibuta, S., Mashimo, T., Ohara, A., Zhang, P., Yoshiya, I., Intracerebroventricular administration of a nitric oxide-releasing compound, NOC-18, produces thermal hyperalgesia in rats, 103
Shinosaki, K., see Inouye, T., 29
Siest, G., see Jolivalt, C., 65
Silverman, W.F., see Yanai, J., 153
Simmons, D.R., Spike, R.C., Todd, A.J., Galanin is contained in GABAergic neurons in the rat spinal dorsal horn, 119
Simon E., see Schmid, H.A., 149
Smith C.U.M., see Armstrong, R.A., 13
Soininen, H., Kosunen, O., Helisalmi, S., Mannermaa, A., Paljärvi, L., Talasniemi, S., Ryynänen, M., Riekkinen Sr, P., A severe loss of choline acetyltransferase in the frontal cortex of Alzheimer patients carrying apolipoprotein $\varepsilon 4$ allele, 79
Sparks, D.L, Liu, H., Gross, D.R., Scheff, S.W., Increased density of cortical apolipoprotein E immunoreactive neurons in rabbit brain after dietary administration of cholesterol, 142
Spiegel, K., see Weibel, L., 83
Spike, R.C., see Simmons, D.R., 119
Strassman, A.M., see Eisenberg, E., 17
Suzuki, T., Maeda, J., Funada, M., Misawa, M., The D_3-receptor agonist (±)-7-hydroxy-*N*,*N*-di-n-propyl-2-aminotetralin (7-OH-DPAT) attenuates morphine-induced hyperlocomotion in mice, 45

Takahashi, T., Time-variant structure of auditory cortex of the macaque monkey, 1
Talasniemi, S., see Soininen, H., 79
Talavera, E., Martínez-Lorenzana, G., León-Olea, M., Sánchez -Alvarez, M., Sánchez-Islas, E., Pellicer, F., Histochemical distribution of NADPH-diaphorase in the cerebral ganglion of the crayfish *Cambarellus montezumae*, 177
Taniyama, K., see Koizumi, S., 123
Tateishi, J., see Nakagawa, Y., 209
Tilgmann, C., see Karhunen, T., 57
Todd, A.J., see Simmons, D.R., 119
Toi, S., see Inouye, T., 29
Tomlinson, D.R., see Garrett, N.E., 201

Ulmanen, I., see Karhunen, T., 57
Uvnäs-Moberg, K., see Agren, G., 49

Valkna, A., Juréus, A., Karelson, E., Zilmer, M., Bartfai, T., Langel, Ü., Differential regulation of adenylate cyclase activity in rat ventral and dorsal hippocampus by rat galanin, 75
Vincent-Viry, M., see Jolivalt, C., 65

Weibel, L., Brandenberger, G., Goichot, B., Spiegel, K., Ehrhart, J., Follenius, M., The circadian thyrotropin rhythm is delayed in regular night workers, 83
Weiss, F., see Richter, R.M., 169
Wevers, A., see Schröder, H., 173
Wilson, L., see Anderson-Beck, R., 25
Winblad, B., see Bonkale, W.L., 5

Yamamoto, C., see Hirata, K., 205
Yanai, J., Silverman, W.F., Shamir, D., An avian model for the reversal of 6-hydroxydopamine induced rotating behaviour by neural grafting, 153
Yoshiya, I., see Shibuta, S., 103

Zhang, P., see Shibuta, S., 103
Zilmer, M., see Valkna, A., 75